THE EFFECT OF CHRONIC HYPERCAPNIA ON OXYGEN AFFINITY AND 2, 3 DIPHOSPHOGLYCERATE AS RELATED TO SUBMARINE EXPOSURE

by

Arthur A. Messier and Karl E. Schaefer, M.D.

SUBMARINE MEDICAL RESEARCH LABORATORY
NAVAL SUBMARINE MEDICAL CENTER REPORT NO. 636

Bureau of Medicine and Surgery, Navy Department Research Work Unit MF12.524.006-9028B.06

Reviewed and Approved by:

Charles F. Bell

Reviewed and Approved by:

Charles F. Gell, M.D., D.Sc. (Med)

Scientific Director

SubMedResLab

J. D. Bloom, CDR MC USN

Officer-in-Charge

SubMedResLab

Approved and Released by:

J./E. Stark, Captain, MC, USN

COMMANDING OFFICER

Naval Submarine Medical Center

This document has been approved for public release and sale; its distribution is unlimited.

SUMMARY PAGE

THE PROBLEM

To better define the pH controlled regulation of oxygen re-

affect 2, 3 DPG levels, follow the pH changes with a time lag.

In vitro the pH dependent changes in 2,3 DPG take place within three hours under optimal conditions 16.

In adaptation to high altitude, a fall in pH due to hypocapnia of hyperventilation occurs together with hypoxia resulting in a simultaneous rise in 2,3 DPG and P50.

Evidence indicating that the rise in pH contributes to the increase in 2,3 DPG in hypoxia has been provided by Gerlach et al. 17 who demonstrated that the increase in 2,3 DPG produced by 24-hour exposure of rats to 11% O₂ was abolished when 5% CO2 was added to the low oxygen gas mixture. In human subject, studied at high altitude, the 2, 3 DPG changes reached one-half of its maximum within six hours and the maximum within 24 hours after arriving at altitude¹⁸. Although the 2,3 DPG changes in chronic hypercapnia are in opposite direction, the time sequence reported here is similar to that observed in altitude adaptation.

While the decrease of 2,3 DPG follows the fall in pH with a time lag during the uncompensated phase of respiratory acidosis, the subsequent increase in 2,3 DPG during the compensated phase of respiratory acidosis (three and seven days) does not exhibit a time lag but closely parallels the rise in pH. This difference in the relationship of pH

here demonstrate clearly that the red cell pH changes are the primary cause of the changes in oxygen affinity.

We have demonstrated in chronic hypercapnia the involvement of two mechanisms, which are pH dependent and are known to influence oxygen affinity: 1) changes in 2, 3 DPG, and 2) changes in red cell cations². The latter findings were obtained in previous work carried out under identical experimental conditions. The correlation coefficients between P50 and 2, 3 DPG were 0.93 and between P50 and red cell cations 0.97. There is obviously a relation between 2,3 DPG changes and red cell cation permeability in chronic hypercapnia. It is not possible to state, at this time, how they are linked together.

The observed findings on 2,3 DPG changes in chronic hypercapnia underline the significance of the pH controlled regulation of oxygen release from hemoglobin which has recently been emphasized by Astrup¹⁹ on the basis of clinical studies.

REFERENCES

- K. E. Schaefer, A. A. Messier, Fed. Proc. 28, 720 (1969).
- 2. K. E. Schaefer, A. A. Messier, C. C. Morgan. Respir. Physiol. (1970) In press.
- 3. H. Sommerkamp, et al. <u>Pflugers</u>
 <u>Arch. ges. Physiol.</u> 272: 591-601,

- 5. R. Benesch and R. Benesch, Biochem. and Biophys. Res. Comm. 26, 162-167 (1967).
- 6. A. Chanutin and R. R. Churnish, Arch. Biochem. Biophys. 121, 96 (1967).
- 7. G. M. Guest and S. Rapoport, Am. J. Diseases Children, 58 1072 (1939).
- S. Rapoport and G. M. Guest,
 J. Biol. Chem. 129, 781 (1939).
- R. E. Benesch, R. Benesch,
 C. I. Yu, <u>Biochemistry</u> 8, 2567,
 (1969).
- R. Benesch and R. E. Benesch, Science 160, 83, (1968).
- 11. G. Gardos, Experientia (Basel) 23, 19 (1967).
- 12. A. S. Keitt, Am. J. Med. 41, 762 (1966).
- P. Hilpert, R. G. Fleischmann,
 D. Kempe and H. Bartels, <u>Am.</u>
 J. Physiol. 205, 337-340 (1963).
- 14. H. Bartels and H. Harms,

 <u>Pfluegers Archiv</u>, 268, 334-365

 (1959).
- 15. H. P. Bartels et al. Am. J. Physiol. 205, 331-336, (1963).
- 16. G. Rooth, H. Sommerkamp and H. Bartels, <u>Clin. Sci.</u> 23, 1-4 (1962).

- 17. E. Gerlach, et al. In: Red Cell

 Metabolism and Function, Ed. G.

 Brewer, Plenum Press, (1970) pp.
 155-174.
- 18. C. Lenfant, et al. In: Red Cell

 Metabolism and Function, Ed. G.

 Brewer, Plenum Press (1970) pp.
 203-212.
- 19. P. Astrup. In: Red Cell Metabolism and Function, Ed. G. Brewer, Plenum Press (1970) pp. 67-79.

Security Classification					
DOCUMENT	CONTROL DATA - R &	k D			
(Security classification of title, body of abstract and i	indexing annotation must be e	ntered when the	overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author)	2a, REPORT SECURITY CLASSIFICATION				
NAVAL SUBMARINE MEDICAL RESEA	Unclassified				
LABORATORY, NAVAL SUBMARINE M	2b. GROUP				
THE EFFECT OF CHRONIC HYPERCA DIPHOSPHOGLYCERATE AS RELATI					
4. DESCRIPTIVE NOTES (Type of report and inclusive datas)					
Interim Report					
5. AUTHOR(S) (First name, middle initial, last name)					
Arthur A. MESSIER and Karl E. SCHA	AEFER				
6. REPORT DATE	78. TOTAL NO. O	PAGES	75. NO. OF REPS		
20 July 1970	5		19		
84. CONTRACT OR GRANT NO.	T NO. 98. ORIGINATOR'S REPORT NUM				
	NSMRL Re	port No. 6	36		
b. PROJECT NO.					
MF12.524.006-9028B					
c.	9b, OTHER REPO	RT NO(S) (Any other numbers that may be assigned			
d.					
10. DISTRIBUTION STATEMENT	·		***		
This document has been approved to	for public release a	nd sale; it	s distribution is		
unlimited.	•	•			
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Naval Submarine Medical Center				
	Box 600 Nav				

The relationship between oxygen affinity and 2, 3 diphosphoglycerate (2,3 DPG) in the red cell has been studied in chronic hypercapnia induced by prolonged exposure of guinea pigs to 15% CO₂ in 21% O₂. Red cell pH fell to a minimum after six hours of exposure and subsequently rose without reaching initial values after seven days of exposure. Both oxygen half-saturation pressure (P_{50}) and the level of 2, 3 DPG of the red cells followed the time course of the pH changes. However, both parameters required 24 hours to reach a minimum, following which they increased steadily and were not different from control values after seven days of exposure.

Groton, Connecticut

DD FORM 1473 (PAGE 1)

UNCLASSIFIED
Security Classification

06340

S/N 0102-014-6600

UNCLASSIFIED

Security Classification LINK A LINK B LINK C ROLE ROLE ROLE WT Oxygen transport Oxygen dissociation curves Chronic hypercapnia Oxygen affinity 2-3 diphosphoglycerate

DD FORM 1473 (BACK)
(PAGE 2)

UNCLASSIFIED

Security Classification